Form Approved REPORT DOCUMENTATION PAGE OMB No. 0704-0188 Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching data sources, Prubic reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate of any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. 3. DATES COVERED (From - To) 1. REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE 01/01/2003-12/30/2005 10/23/2007 Final 4. TITLE AND SUBTITLE 5a. CONTRACT NUMBERS Structured and Inhibited Mixing on the Continental shelf **5b. GRANT NUMBER** N00014-03-1-0335 **5c. PROGRAM ELEMENT NUMBER** 5d. PROJECT NUMBER 6. AUTHOR(S) 133335SP Duda, Timothy F. 5e. TASK NUMBER 5f. WORK UNIT NUMBER 8. PERFORMING ORGANIZATION REPORT 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NUMBER Woods Hole Oceanographic Institution 266 Woods Hole Rd. Woods Hole, MA 02543 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSORING/MONITORING ACRONYM(S) Office of Naval Research One Liberty Center 11. SPONSORING/MONITORING AGENCY 875 North Randolph Street, Suite 1425 REPORT NUMBER Arlington, VA 22203 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited 13. SUPPLEMENTARY NOTES 14. ABSTRACT A goal is to understand diapycnal mixing process during the stratified summer season in the temperate continental shelf environment. This understanding is required to correctly account for the effects of mixing in coastal ocean models. These effects influence modeled (and actual) density fields, and thus flows, as well as modeled and actual distributions of dissolved and suspended material.

17. LIMITATION OF

ABSTRACT

c. THIS PAGE

18. NUMBER

OF PAGES

4

15. SUBJECT TERMS

a. REPORT

16. SECURITY CLASSIFICATION OF:

b. ABSTRACT

Standard Form 298 (Rev. 8-98) Prescribed by ANSI-Std. Z39-18

19a. NAME OF RESPONSIBLE PERSON

19 b. TELEPHONE NUMBER (Include are code)

Timothy F. Duda

508-289-2495

Structured and Inhibited Mixing on the Continental Shelf

Timothy F. Duda
Applied Ocean Physics and Engineering Department, MS 11
Woods Hole Oceanographic Institution, Woods Hole, MA 02543
phone: (508) 289-2495 fax: (508) 457-2194 email: tduda@whoi.edu

Award Number: N00014-03-1-0335

Final Project Report

LONG TERM GOALS

A goal is to understand diapycnal mixing process during the stratified summer season in the temperate continental shelf environment. This understanding is required to correctly account for the effects of mixing in coastal ocean models. These effects influence modeled (and actual) density fields, and thus flows, as well as modeled and actual distributions of dissolved and suspended material.

OBJECTIVES

The objective of this project was to determine whether intermittent diapycnal mixing events (i.e. turbulent events that cause cross-isopycnal mass flux) are sensitive to the background conditions in a specific way. The hypothesis is that, within a field exhibiting a step-like density structure, mixing will be more rapid in less-stratified layers than in more stratified interfaces. Such an effect would cause density-gradient perturbations to grow, an effect sometimes known as the Phillips instability (Phillips, 1972; see also Balmforth et al 1998). This behavior is analogous to a negative eddy diffusivity K. Evidence of gradient-dependent K was found in the ONR Coastal Mixing and Optics (CMO) program dye diffusion studies, and in towed-microscale conductivity measurements obtained in conjunction with the dye studies (Duda and Rehmann, 2002). We wish to expand on the CMO work by obtaining and using more data of better quality than previously available, confirming and/or improving the result.

APPROACH

An extensive set of microscale conductivity data and supporting data were obtained with a towed sensor package (called Micro-Tow, MT) during a seven-day research cruise on RV Endeavor during August 2004. The towed platform and the onboard data-acquisition electronics were of our own design.

The measured conductivity gradient variance, which should be proportional to temperature gradient variance and to χ , the thermal dissipation rate, can be compared to the vertical temperature gradient, with both estimated over short time intervals. A definitive correlation of the two quantities, and a relationship with a smaller exponent than χ proportional to dT/dz

to the first power would indicate eddy diffusivity K inversely dependent on the gradient, supporting the Phillips instability.

WORK COMPLETED

The field program was carried out August 8-14, 2004. This was RV Endeavor cruise EN396, sailing from/to Narragansett, RI. This ship is operated by the University of Rhode Island. All systems functioned well during the cruise. The MT system was in the water collecting data for approximately 92 hours, representing 60% of the time at sea (Figure 1). For an additional 27 hours we collected data with the University of Rhode Island automatic undulating Scanfish system, which we specially equipped with microstructure sensors through a collaboration with the Naval Research Laboratory (Dr. David Walsh).

We had partial success at staying in water with a desirable near-zero salinity gradient. We believe that for most of the time we remained in water with small enough salinity gradient to require only small correction factors in our analysis.

Prior to the field work, the existing towed conductivity vehicle was upgraded to carry a second microstructure sensor. Also, new ship-board data acquisition and data-visualization reporting software was written.

RESULTS

Data acquisition results were excellent. All towed systems and ship-board systems worked well for the duration of the experiment. Most if not all of the data appear to be usable and of excellent quality. Figure 2 shows example temperature, salinity, and conductivity gradient variance from one transect.

There were two impediments that prevented finalization of a publication of results in a journal article during the course of the project. First, we had trouble successfully locating water with low salinity gradient and with doubly-diffusively stable stratification, which would ensure lowest possible analysis errors and the smallest error bars. Thus, extra work is required to improve statistical reliability beyond that of our existing publication. The data must be culled and gone over in additional ways to extract the best data and compute a new χ - to - vertical temperature gradient correlation result. Next, loss of communication synchronization between submerged vehicle and ship-board PC caused anomalies in the data stream which required extra time to work around. Thus, final correlations and investigation were not obtained.

TRANSITIONS

The tow platform built by us for this type of work has been used by Naval Research Laboratory personnel in conjunction with monostatic high-frequency flow-visualization acoustics to study nonlinear internal gravity waves in the South China Sea. In addition, NRL has built a sensor package to make towed microscale measurements, similar to our measurements, in order to study mixing processes in coastal and near-surface waters. This is the sensor package that was appended to the auto-undulating Scanfish during our cruise.

REFERENCES

Balmforth, N. J., S. G. Llewellyn Smith, and W.R. Young, Dynamics of interfaces and layers in a stratified turbulent fluid, *J. Fluid Mech.*, 355, 329-358, 1998.

Duda, T. F., and C. R. Rehmann, Systematic microstructure variability in double-diffusively stable coastal waters of nonuniform density gradient. J. Geophys. Res., 107(C10), 3145, doi:10.1029/2001JC000844, 2002.

Phillips, O. M., Turbulence in a strongly stratified fluid – is it unstable?, *Deep-Sea Res.*, 19, 79-81, 1972.




Figure 1: Track lines of Endeavor along which MT data were collected on the continental shelf south of New England.